



## Sources and proxy potential of long chain alkyl diols in lacustrine environments

Sebastiaan W. Rampen<sup>a,\*</sup>, Mariska Datema<sup>b</sup>, Marta Rodrigo-Gámiz<sup>a</sup>,  
Stefan Schouten<sup>a,b</sup>, Gert-Jan Reichart<sup>b,c</sup>, Jaap S. Sinninghe Damsté<sup>a,b</sup>

<sup>a</sup> NIOZ Royal Netherlands Institute for Sea Research, Department of Marine Organic Biogeochemistry, P.O. Box 59, 1790 AB Den Burg, Texel, The Netherlands

<sup>b</sup> Utrecht University, Faculty of Geosciences, Department of Earth Sciences, P.O. Box 80.021, 3508 TA Utrecht, The Netherlands

<sup>c</sup> NIOZ Royal Netherlands Institute for Sea Research, Department of Marine Geology and Chemical Oceanography, P.O. Box 59, 1790 AB Den Burg, Texel, The Netherlands

Received 10 May 2014; accepted in revised form 27 August 2014; available online 16 September 2014

### Abstract

Long chain 1,13- and 1,15-alkyl diols form the base of a number of recently proposed proxies used for climate reconstruction. However, the sources of these lipids and environmental controls on their distribution are still poorly constrained. We have analyzed the long chain alkyl diol (LCD) composition of cultures of ten eustigmatophyte species, with three species from different families grown at various temperatures, to identify the effect of species composition and growth temperature on the LCD distribution. The results were compared with the LCD distribution of sixty-two lake surface sediments, and with previously reported LCD distributions from marine environments. The different families within the Eustigmatophyceae show distinct LCD patterns, with the freshwater family Eustigmataceae most closely resembling LCD distributions in both marine and lake environments. Unlike the other two eustigmatophyte families analyzed (Monodopsidaceae and Goniochloridaceae), C<sub>28</sub> and C<sub>30</sub> 1,13-alkyl diols and C<sub>30</sub> and C<sub>32</sub> 1,15-alkyl diols are all relatively abundant in the family Eustigmataceae, while the mono-unsaturated C<sub>32</sub> 1,15-alkyl diol was below detection limit. In contrast to the marine environment, LCD distributions in lakes did not show a clear relationship with temperature. The Long chain Diol Index (LDI), a proxy previously proposed for sea surface temperature reconstruction, showed a relatively weak correlation ( $R^2 = 0.33$ ) with mean annual air temperature used as an approximation for annual mean surface temperature of the lakes. A much-improved correlation ( $R^2 = 0.74$ ,  $p$ -value  $< 0.001$ ) was observed applying a multiple linear regression analysis between LCD distributions and lake temperatures reconstructed using branched tetraether lipid distributions. The obtained regression model provides good estimates of temperatures for cultures of the family Eustigmataceae, suggesting that algae belonging to this family have an important role as a source for LCDs in lacustrine environments, or, alternatively, that the main sources of LCDs are similarly affected by temperature as the Eustigmataceae. The results suggest that LCDs may have the potential to be applicable as a palaeotemperature proxy for lacustrine environments, although further calibration work is still required.

© 2014 Elsevier Ltd. All rights reserved.

### 1. INTRODUCTION

After the first discovery of long chain alkyl diols (LCD) in Black Sea sediments (De Leeuw et al., 1981), it has

become clear that they are widely distributed in both marine and freshwater environments (Versteegh et al., 1997). LCDs occur as several structurally related series, and are derived from distinct sources. The marine diatom genus *Proboscia*, for example, has been identified as a major source for both saturated and mono-unsaturated C<sub>28</sub> and

\* Corresponding author.

E-mail address: [sebastiaan.rampen@nioz.nl](mailto:sebastiaan.rampen@nioz.nl) (S.W. Rampen).

$C_{30}$  1,14-alkyl diols (Sinninghe Damsté et al., 2003), whereas the freshwater fern *Azolla* has been linked to  $C_{30}$ – $C_{36}$  1,ω20-alkyl diols (Speelman et al., 2009). However, the major sources of the most common long chain diols in both marine and freshwater environments, i.e.  $C_{28}$ – $C_{32}$  1,13- and 1,15-alkyl diols (also indicated as 1,ω18- and 1,ω16-alkyl diols; Versteegh et al., 1997), have not been unambiguously identified yet. Eustigmatophyte algae are the only organisms known to produce both 1,13- and 1,15-alkyl diols (Volkman et al., 1992, 1999; Méjanelle et al., 2003; Shimokawara et al., 2010) but their importance as a source for LCDs, particularly in marine environments, is uncertain. These algae are rarely reported in the marine environment and the LCD distribution of cultured marine eustigmatophytes is generally dominated by saturated and mono-unsaturated  $C_{32}$  1,15-alkyl diols (Volkman et al., 1992, 1999), in contrast to the marine environments, where the mono-unsaturated  $C_{32}$  1,15-alkyl diol is not detected and the saturated  $C_{32}$  1,15-alkyl diol is present in relatively low abundances (~0–30% of all 1,13- and 1,15-alkyl diols; Rampen et al., 2012). For lake environments, the major sources of these LCDs are also uncertain. The similarity of the LCD distribution of a *Nannochloropsis* sp. and Lake Baikal sediments suggests Eustigmatophyceae as a possible source in lake environments (Shimokawara et al., 2010). In addition, Villanueva et al. (2014) observed similar patterns between 18S rRNA gene copy numbers of Eustigmatophyceae and the concentration of LCDs in the water column of the stratified Lake Challa (East Africa). The results from that study suggested that not-yet described Eustigmatophyte groups are the major producers of the LCDs. The occurrence of uncultivated Eustigmatophyte groups is in agreement with a study on the phylogeny of Eustigmatophyceae isolates obtained from lakes, ponds and bogs from Itasca State Park, revealing the existence of several new genera and species (Prior et al., 2009), indicating that our knowledge on this algal group is still fairly limited.

LCDs are increasingly used as palaeoenvironmental indicators. In 1997, Versteegh et al. proposed the diol index, a ratio between  $C_{30}$  and  $C_{32}$  1,15-alkyl diols, as a marker for upwelling or freshwater influence in marine environments. More recently, ratios between 1,14-alkyl diols and 1,13- or 1,15-alkyl diols have also been applied as indicators for upwelling (Rampen et al., 2008; Willmott et al., 2010; Seki et al., 2012), while 1,ω20-alkyl diols have been proposed as markers for the past occurrence of *Azolla* and, consequently, freshwater conditions (Speelman et al., 2009). Furthermore, a ratio between LCDs and tetrahymanol (Romero-Viana et al., 2012), and the deuterium isotopic composition of the  $C_{32}$  1,15-alkyl diol (Romero-Viana et al., 2013) have been introduced as indicators for rainfall variability. Rampen et al. (2012) introduced the Long chain Diol Index (LDI), based on a ratio between  $C_{30}$  1,15-alkyl diol and  $C_{28}$  and  $C_{30}$  1,13-alkyl diols, as a proxy for the reconstruction of past sea surface temperature. The LDI has subsequently been used in paleoceanographic studies and shows sea temperature reconstructions that are similar to those obtained with other organic proxies (Rampen et al., 2012; Lopes dos Santos et al., 2013; Rodrigo-Gámiz et al., 2014). Consequently,

as most of these LCD proxies involve  $C_{28}$ – $C_{32}$  1,13- and 1,15-alkyl diols, it becomes increasingly relevant to identify their sources in both marine and lacustrine environments.

In order to shed further light on the role of eustigmatophyte algae as a potential source for LCDs, we analyzed LCD distributions in ten different species, selected from all phylogenetic branches within the Eustigmatophyceae for which algal cultures were available. Furthermore, three species from different groups were grown at a range of temperatures to determine how growth temperature affects the LCD composition. The culture results obtained were compared with published data from the marine natural environment (Rampen et al., 2012). Furthermore, we analyzed LCD distributions in surface sediments from sixty-two lakes, previously used for a study on branched glycerol dialkyl glycerol tetraethers (branched GDGTs; Blaga et al., 2010). This lacustrine LCD dataset was compared with the LCD distributions from the cultures and the marine dataset, and finally used to test whether LCDs could also be utilized as lake temperature proxies.

## 2. MATERIALS AND METHODS

Unialgal cultures of ten eustigmatophyte species were grown in batch cultures with an appropriate culture medium, using a 12/12 h light/dark cycle, and harvested at the end of the log growth phase (Table 1 and Supplementary data). The cultures of *Goniochloris sculpta* grown at 8 and 14 °C, which did not grow well and never reached the exponential growth phase, were harvested after 49 days. Algal material was collected either by filtration on pre-combusted Whatman GF/C 47 mm filters or centrifugation (Table 1). The samples were saponified according to Rampen et al. (2011) and the extracts obtained were fractionated into apolar and polar fractions using a glass pipette column filled with activated alumina and elution with hexane/dichloromethane (9/1; vol./vol.) and dichloromethane/methanol (1/1; vol./vol.), respectively. Aliquots of the polar fractions of *Nannochloropsis gaditana*, grown at 14 °C, and *G. sculpta*, grown at 16 °C were dissolved in hexane and derivatized with dimethyl-disulfide (DMDS) (activated with iodine in diethyl ether at 40 °C overnight) to determine the position of the double bond (Buser et al., 1983). Prior to analysis by gas chromatography/mass spectrometry (GC/MS), polar fractions were silylated by adding *N,O*-bis(trimethylsilyl)trifluoroacetamide (BSTFA) and pyridine and heating the mixture in an oven at 60 °C for 20 min. GC/MS analyses were performed using a Thermofinnigan TRACE gas chromatograph equipped with a fused silica column (25 m × 0.32 mm) coated with CP Sil-5 (film thickness 0.12 μm) and He as carrier gas, using GC conditions as described by Rampen et al. (2008). The gas chromatograph was coupled to a Thermofinnigan DSQ quadrupole mass spectrometer scanning a mass range of *m/z* 50–800 at 3 scans per second and operating at an ionization energy of 70 eV. LCDs were quantified using the relative intensities of characteristic fragments in their mass spectra (cf. Versteegh et al., 1997). 18S rRNA gene sequences from the same algal species as analyzed in this study were obtained from Genbank (<http://www.ncbi.nlm.nih.gov/>)

Table 1  
Eustigmatophyte culture information.

Culture	Original environment	Medium <sup>a</sup>	Temperature (°C)	Duration (days)	Harvest method	18S rRNA gene sequence
<i>Monodopsidaceae</i>						
<i>Nannochloropsis gaditana</i> (CCAP 849/5)	Marine	F/2	8,14,16, 20	38, 32, 31, 35	Filter	AF045036
<i>Nannochloropsis oceanica</i> (CCAP 849/10)	Marine	F/2	15	9	Filter	JF489982
<i>Monodopsis subterranea</i> (CCAP 848/1)	Freshwater	3N BBM + V	20	35	Filter	U41054
<i>Eustigmataceae</i>						
<i>Vischeria punctata</i> (CCAP 887/3)	Soil	3N BBM + V	20	50	Filter	FJ858972
<i>Vischeria stellata</i> (CCAP 887/4)	Soil	3N BBM + V	8,14,16, 20	123, 56, 56, 29	Filter	HQ710570
<i>Chloridella simplex</i> (SAG 51.91)	Snow	JM	16	87	Filter	KF848923
<i>Eustigmator vischeri</i> (CCAP 860/7)	Soil	3N BBM + V	20	41	Filter	FJ858973
<i>Goniocloridaceae</i>						
<i>Goniocloris sculpta</i> (SAG 29.96)	Freshwater	JM	8, 14,16, 19.5	68, 21, 49, 49	Centrifuge	FJ858970
<i>Microtalis aquatica</i> (CCMP 3153)	Freshwater	MDY	20	35	Filter	HQ710564
<i>Pseudostaurastrum enorme</i> (SAG 11.85)	Freshwater	Modified DesMid	16	53	Filter	EF044312

<sup>a</sup> Detailed information on culture media can be obtained from <http://www.ccap.ac.uk/>.

genbank/, Table 1) and a Neighbor Joining phylogenetic tree was created using distances calculated according to Kimura (1980) and clustering described by Saitou and Nei (1987), which was transformed into an ultrametric tree using the PATHd8 program (Britton et al., 2007).

Sixty-two lake surface sediments were analyzed for their LCD content. Sediments had been extracted previously and branched GDGT compositions have been reported by Blaga et al. (2010). All polar fractions were silylated as described above, and analyzed for LCD distributions using an Agilent Technologies 7890A GC system equipped with a fused silica column (30 m × 0.32 mm) coated with HP-5MS (film thickness 0.25 μm) and He as carrier gas. The gas chromatograph was coupled to an Agilent Technologies 5975C VL MSD operating with an ionization energy of 70 eV using GC conditions as described by Rampen et al. (2008). LCDs were identified using a full scan method, scanning a mass range of 50–800 and quantified using a SIM method, using characteristic fragments in the mass spectra, i.e. *m/z* 299, 313, 327, 341 and 355 (cf. Rampen et al., 2008, 2012). LCD distributions were correlated with annual mean air temperatures obtained from nearby climate observation stations (KNMI, 1998) instead of temperatures provided by Blaga et al. (2010), following the approach of Pearson et al. (2011). SigmaPlot 12.0 was used for regression analyses, while analyses of covariance were performed using SYSTAT 13.

### 3. RESULTS

#### 3.1. LCDs in eustigmatophyte cultures

For this study we analyzed LCD distributions in cultures of ten eustigmatophyte species, classified within the families Eustigmataceae, Monodopsidaceae and Goniocloridaceae (classification according to Fawley et al., in press, and references cited therein). Although the major environmentally significant groups of eustigmatophytes

are probably not yet available in culture (Villanueva et al., 2014), we attempted to obtain the broadest phylogenetic variety of eustigmatophyte species currently possible. From each of the three available eustigmatophyte families, one species was grown at four different temperatures (8, 14, 16 and 20 °C; see Table 1).

All eustigmatophyte species analyzed were found to contain LCDs (long chain diol composition of the eustigmatophyte species are available online in Supplementary Table 1 and at <http://pangaea.de>). The LCD distributions of species of the family Monodopsidaceae were dominated by the C<sub>32</sub> 1,15-alkyl diol, and contained lower fractional abundances (relative to the sum of all LCDs) of C<sub>30</sub> and C<sub>31</sub> 1,15-alkyl diols and the C<sub>30</sub> 1,13-alkyl diol (Fig. 1). The two marine species in this study, *N. gaditana* and *N. oceanica*, contained relatively high fractional abundances of the mono-unsaturated (i.e. with the double bond at C-27) C<sub>32</sub> 1,15-alkyl diol. *N. oceanica* also contained a mono-unsaturated C<sub>32</sub> 1,17-alkyl diol, but unsaturated LCDs were not detected in *Monodopsis subterranea*.

*G. sculpta* and *Microtalis aquatica*, both belonging to the Goniocloridaceae, showed an LCD distribution similar to those of the studied species of the Monodopsidaceae, with predominantly saturated and mono-unsaturated C<sub>32</sub> 1,15-alkyl diols, together with C<sub>30</sub> and C<sub>31</sub> 1,15-alkyl diols (Fig. 1). The double bond position for *G. sculpta*, grown at 16 °C, was determined to be at C-27 as observed for the Monodopsidaceae. However, unlike the Monodopsidaceae, *G. sculpta* and *M. aquatica* contained only 1,15-alkyl diols and no C<sub>30</sub> 1,13-alkyl diol. *Pseudostaurastrum enorme*, classified in the same family as *G. sculpta* and *M. aquatica*, had a substantially different LCD distribution, comprising saturated and mono-unsaturated C<sub>32</sub> 1,15-alkyl diols, but also C<sub>28</sub> and C<sub>30</sub> 1,13-, 1,14- and 1,15-alkyl diols and only a low contribution of C<sub>31</sub> 1,15-alkyl diol (<1%).

Eustigmataceae showed a distinct LCD distribution, dominated by the C<sub>28</sub> 1,13-alkyl diol, C<sub>30</sub> 1,15-alkyl diol and C<sub>32</sub> 1,15-alkyl diol, together with lower abundances




Fig. 1. LCD carbon number distributions (expressed as fractional abundances) of the cultures of the ten studied species of Eustigmatophyceae. Three species were grown at four different temperatures as indicated. For each carbon number the various LCD isomers are indicated by a colour code. For reference, the molecular phylogeny of the Eustigmatophyceae based on the 18S rRNA gene, is provided.

of  $C_{29}$  and  $C_{30}$  1,13-alkyl diols and  $C_{31}$  1,15-alkyl diol (Fig. 1). In addition, *Vischeria stellata* contained a mixture of  $C_{34}$  alkyl diols, dominated by the 1,19-diol isomer. A mono-unsaturated  $C_{32}$  1,15-alkyl diol was not detected in any of the Eustigmataceae cultures.

The LCD distributions of the three different families do not substantially change with growth temperature, with the exception of a decrease in the fractional abundance of the monounsaturated  $C_{32}$  1,15-alkyl diol compared to the

saturated  $C_{32}$  1,15-alkyl diol with increasing growth temperature as observed for both *N. gaditana* and *G. sculpta* (Figs. 1 and 2f).

### 3.2. LCDs in lake sediments

We also analyzed the LCD composition of surface sediments of sixty-two lakes, mainly from Europe and South America (Blaga et al., 2010). LCD concentrations in these




Fig. 2. Fractional abundances (F) of individual LCDs (a–d; C<sub>28</sub> 1,13-alkyl diol, C<sub>30</sub> 1,15-alkyl diol, C<sub>30</sub> 1,13-alkyl diol, C<sub>32</sub> 1,15-alkyl diol, respectively), the LDI ratio (e) and the fractional abundances of unsaturated LCDs (f) of Eustigmatophyceae cultures plotted versus growth temperature. Data of Rampen et al. (2009) for *Proboscia* diatoms is also plotted in (f).

lakes were generally low, requiring the use of a selective ion monitoring (SIM) method (Rampen et al., 2012). Even when using SIM, LCDs were below detection limit in four lake sediments. The remaining fifty-eight lakes show significant variation in their LCD composition (Fig. 3; Supplementary Table 2). LCDs predominantly consist of C<sub>28</sub>,

C<sub>30</sub> and C<sub>32</sub> chain lengths (Fig. 3a). On average, C<sub>30</sub> alkyl diols were most abundant in the studied lake sediments, contributing 20–50% of all LCDs and being the most dominant LCDs in 50% of all lakes. However, in the South American lakes, C<sub>32</sub> alkyl diols were the most dominant LCDs in 56% of all lakes investigated, while C<sub>28</sub> alkyl diols




Fig. 3. Bar plots showing the chain length distributions (a) and distributions of the different LCD isomers with chain lengths of 28–32 (b–f, respectively) carbon atoms for the eustigmatophyte cultures and the lake sediments.

were relatively more abundant in the European lakes compared to South American lakes. Most lakes also contained small amounts of  $C_{34}$  alkyl diols, but these were not quantified.

The  $C_{32}$  1,15-alkyl diol was present in all of the lake sediments, generally contributing >10% of all LCDs, and the most abundant LCD in almost 75% of all South American lakes and 30% of all European lakes.  $C_{30}$  1,13- and

1,15-alkyl diols were also present in all lakes, together with relatively low  $C_{30}$  1,14- and 1,16-alkyl diols. In the European lakes,  $C_{28}$  alkyl diols mainly consisted of the 1,13-alkyl diol, which was the most dominant LCD in 50% of these lakes, while  $C_{28}$  1,13- and 1,14-alkyl diol fractional abundances in the South American lakes were similar, but low. The mono-unsaturated  $C_{32}$  1,15-alkyl diol was only present in low abundances or below detection



limit (not shown). Concentrations were too low to determine the exact position of the double bond, but mass spectral fragmentation pattern of this sedimentary LCD indicated that the double bond was positioned between C<sub>16</sub> and C<sub>32</sub>.

## 4. DISCUSSION

### 4.1. Comparison of LCD distributions in cultures, and lake and marine sediments

C<sub>28</sub> and C<sub>30</sub> 1,13-alkyl diols and C<sub>30</sub> and C<sub>32</sub> 1,15-alkyl diols were the dominant LCDs in Eustigmatophyceae cultures, but the three phylogenetic clusters of Eustigmatophyceae did show clear differences in their LCD patterns (Figs. 1–3). Most obvious is the presence of the mono-unsaturated C<sub>32</sub> 1,15-alkyl diol in Monodopsidaceae and Goniochloridaceae and its absence in the Eustigmataceae. It is also remarkable that 1,13-alkyl diols were not detected in the *Goniochloris* and *Microtalis* species, and species of the family Monodopsidaceae contained relatively low amounts of the C<sub>30</sub> 1,13-alkyl diol, while Eustigmataceae and the species *Pseudostaurastrum enorme* contained both C<sub>28</sub> and C<sub>30</sub> 1,13-alkyl diols. The species *P. enorme* differed from all other studied algae in containing relatively high amounts of 1,14-alkyl diols (>30% of the total LCD composition). Long chain alkyl diol distributions in Monodopsidaceae and Eustigmataceae are comparable to those previously reported by Volkman et al. (1992, 1999), Méjanelle et al. (2003) and Shimokawara et al. (2010), while this is the first report on long chain alkyl diol distributions of Goniochloridaceae. It was not possible to grow all cultures in the same culture medium, and the effect of the different culture media (Table 1) on the LCD distributions remains untested. However, differences between phylogenetically related species, grown in different media, are relatively small, while species, belonging to different clusters but grown in the same culture media, show substantially different LCD distributions.

Most (i.e. >90%) of the lake sediments studied contained LCDs, indicating that these lipids are common compounds in lacustrine environments. LCDs in lakes predominantly comprise C<sub>28</sub> and C<sub>30</sub> 1,13-alkyl diols and C<sub>30</sub> and C<sub>32</sub> 1,15-alkyl diols (Fig. 3), as observed in marine environments (Versteegh et al., 1997; Rampen et al., 2012). The relative abundances of the three most abundant LCDs, the C<sub>28</sub> 1,13-alkyl diol, C<sub>30</sub> 1,15-alkyl diol and C<sub>32</sub> 1,15-alkyl diol, are plotted in a ternary diagram to compare LCD distributions from the different eustigmatophyte families, marine sediments and lake sediments (Fig. 4). This shows that marine surface sediments generally have different LCD distributions compared to the lake sediments and the studied eustigmatophyte species. In marine surface sediments, the C<sub>32</sub> 1,15-diol rarely contributes >30% to the sum of these three LCDs, while this LCD is often the dominant alkyl diol in lake surface sediments and Eustigmatophyceae. It is remarkable that four of the marine sediments containing >30% C<sub>32</sub> 1,15-alkyl diol are from the Hudson Bay, a large inland sea strongly influenced by riverine input (e.g. Déry et al., 2011). Indeed, it has already been noted that the abundance of C<sub>32</sub> 1,15-alkyl diol is higher in brackish and

freshwater areas compared to the marine environment (Versteegh et al., 1997), suggesting that a high abundance of this LCD may signify freshwater/brackish conditions. However, a substantial number of freshwater lakes are characterized by a relatively low C<sub>32</sub> 1,15-alkyl diol abundance, similar to those in the marine environment (Fig. 4) indicating that a low relative abundance of this LCD may not in itself signify marine conditions. It remains uncertain what determines the C<sub>32</sub> 1,15-alkyl diol abundance; no significant correlations were observed between C<sub>32</sub> 1,15-alkyl diol abundances and lake size or depth, or pH (data not shown), but the different LCD distributions of the different eustigmatophyte families (Fig. 1) suggest that this could be species-related.

LCD distributions in species belonging to the family Eustigmataceae are, compared to the other families, relatively similar to those in lake and marine sediments (Fig. 4). In contrast, species of the families Monodopsidaceae and Goniochloridaceae contain high fractional abundances of the mono-unsaturated C<sub>32</sub> 1,15-alkyl diol, which is not detected or present at low relative abundances in marine and lacustrine environments. Possibly, the mono-unsaturated C<sub>32</sub> 1,15-alkyl diol may be preferentially degraded compared to the saturated LCDs, or transformed into saturated LCDs, potentially explaining this discrepancy. However, the mono-unsaturated C<sub>32</sub> 1,15-alkyl diol is also absent or present in low abundance in suspended particulate matter and descending particles in marine and lacustrine environments (e.g. Rampen et al., 2007, 2008; Villanueva et al., 2014), suggesting that the Monodopsidaceae and Goniochloridaceae species are not major sources for LCDs in aquatic environments. Although their LCD distributions indicate that species of the family Eustigmataceae could be producers of these lipids in aquatic environments, a recent gene-based study by Villanueva et al. (2014) revealed that most of the Eustigmatophyceae clones in the East African Lake Challa belonged to novel Eustigmatophyceae groups, suggesting that the major sources for LCDs in the environment may not yet be available in culture.

### 4.2. Impact of temperature on LCD distributions

*N. gaditana* and *G. sculpta*, grown at temperatures between 8 and 20 °C, show a decrease in the fractional abundance of mono-unsaturated C<sub>32</sub> 1,15-alkyl diol, while the fractional abundance of the saturated C<sub>32</sub> 1,15-alkyl diol increased with increasing temperature (Fig. 2). A similar increase in fractional abundance of unsaturated long chain 1,14-alkyl diols with increasing temperature was observed for *Proboscia* diatoms (Rampen et al., 2009). Changing the degree of unsaturation is a well-known environmental adaptation mechanism, often used by organisms to maintain a constant lipid fluidity and membrane permeability (Russell and Fukunaga, 1990; Suutari and Laakso, 1994). However, as discussed above, in both marine (Versteegh et al., 1997; Rampen et al., 2007, 2008) and lake sediments (Villanueva et al., 2014; this study), mono-unsaturated C<sub>32</sub> 1,15-alkyl diols are rarely abundant. Thus, in aquatic environments, probably Eustigmatophyceae species other than Monodopsidaceae and Goniochloridaceae are




Fig. 4. Ternary diagram showing relative abundances of  $C_{28}$  1,13-alkyl diol,  $C_{30}$  1,15-alkyl diol and  $C_{32}$  1,15-alkyl diol from cultures (this study), lake surface sediments (this study) and marine surface sediments (data published by Rampen et al., 2012). The green line indicates 30%  $C_{32}$  1,15-alkyl diol, which is a maximum value for most marine sediments.

the major sources for LCDs. These species may have adopted other methods to adapt their lipid composition to growth temperature. In marine surface sediments, the relative abundance of both  $C_{28}$  and  $C_{30}$  1,13-alkyl diols decrease, and that of the  $C_{30}$  1,15-alkyl diol increases with increasing temperature, which forms the basis of the LDI paleotemperature proxy (Rampen et al., 2012). LDI-values for the *G. sculpta* cultures grown at different temperatures are 1 due to the absence of 1,13-alkyl diols, while LDI values for *N. gaditana* cultures show a negative trend with increasing growth temperature (Fig. 2e). Eustigmataceae do show similar temperature trends for the fractional abundance of individual LCDs and the LDI as those observed for marine sediments (Fig. 2), but with significant offsets, which may be due to different growth conditions (e.g. salinity as the Eustigmataceae species cultivated are freshwater species). Alternatively, it could suggest that species of this family are not a major source for LCDs in the marine environment.

We also investigated the influence of temperature on the distribution of LCDs in lake sediments, by plotting relative abundances (i.e. the abundance of an individual LCD divided by the summed abundances of  $C_{28}$  and  $C_{30}$  1,13-alkyl diols and  $C_{30}$  and  $C_{32}$  1,15-alkyl diols) of individual 1,13- and 1,15-alkyl diols in the lake surface sediments versus temperature. Preferably, measured in-situ lake temperatures should be used but these were often not available for the lakes studied. As an alternative, mean annual air

temperatures (MAAT) obtained from nearby weather stations were used (KNMI, 1998), as lake water temperatures are often strongly correlated with MAAT (Livingstone and Lotter, 1998). Only weak or no correlations are observed for the relationship between relative LCD abundances and temperature (Fig. 5a–d), which is in contrast to correlations observed for the relative abundances of  $C_{28}$  and  $C_{30}$  1,13-alkyl diols and  $C_{30}$  1,15-alkyl diols with sea surface temperature in marine surface sediments (Rampen et al., 2012). Nevertheless, LDI values of the LCDs in the lake sediments do show a weak but significant correlation with MAAT ( $R^2 = 0.33$ ,  $p$ -value  $< 0.001$ ,  $n = 56$ ; Fig. 5e). For branched GDGTs it has been shown that improved correlations may be obtained between branched GDGT relative abundances and MAAT when multiple linear regression is used (Tierney et al., 2010; Loomis et al., 2012). Therefore, we applied a multiple linear regression on the relative abundances of the four most dominant LCDs, e.g.  $C_{28}$  and  $C_{30}$  1,13-alkyl diol,  $C_{30}$  and  $C_{32}$  1,15-alkyl diol and MAAT. The following model was obtained (Fig. 5f):

$$\begin{aligned} \text{Temperature} (^{\circ}\text{C}) = & 24.6 - 32.0 * \text{RA}_{C_{28}1,13\text{-alkyl diol}} \\ & - 59.0 * \text{RA}_{C_{30}1,13\text{-alkyl diol}} - 7.4 \\ & * \text{RA}_{C_{30}1,15\text{-alkyl diol}} \end{aligned} \quad (1)$$

where RA indicates the relative abundance of an individual LCD. Because of the closed sum effect, the relative abundance of the  $C_{32}$  1,15 alkyl diol is not expressed in






Fig. 5. Cross plots of the relative abundances (RA) of C<sub>28</sub> 1,13-alkyl diol, C<sub>30</sub> 1,15-alkyl diol, C<sub>30</sub> 1,13-alkyl diol, C<sub>32</sub> 1,15-alkyl diol vs. mean annual air temperatures (MAAT) (a–d, respectively), LDI values vs. MAAT (e) and (f) MAAT vs. LCD inferred temperatures (Eq. 1), obtained from lake sediments. Light blue lines indicate the relationships between the fractional abundance of LCDs (a–d) and LDI (e) in marine surface sediments and sea surface temperature as reported by Rampen et al. (2012). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

formula (1) but does affect the outcome of the equation. The correlation observed for this model ( $R^2 = 0.53$ ,  $p$ -value  $< 0.001$ ,  $n = 56$ ; Fig. 5f) is stronger than observed for the LDI, but still much weaker than that observed between the LDI and sea surface temperature for the marine environment (i.e.,  $R^2 = 0.97$ , Rampen et al., 2012). Part of the scatter may be due to inaccuracies in the lake temperature data used, due to differences in location and altitude of the lakes and the weather stations, but

the correlation did not improve after correcting for altitude differences using lapse rates of 6.5 or 5.1 °C km<sup>-1</sup> (Stone and Carlson, 1979). Another reason for the observed scatter may be that LCDs are produced in a specific time period during the annual cycle. Other proxies are known to be affected by seasonal production maxima (e.g. Castañeda and Schouten, 2011 and references cited therein). However, we also did not observe a significant improvement of the correlation between LCDs and




Fig. 6. Cross plots of the relative abundances of  $C_{28}$  1,13-alkyl diol,  $C_{30}$  1,15-alkyl diol,  $C_{30}$  1,13-alkyl diol,  $C_{32}$  1,15-alkyl diol (a–d, respectively), and LDI values (e) obtained from cultures and lake sediments vs. temperature and branched GDGT-derived summer temperatures, respectively, and (f) temperature and branched GDGT-derived summer temperatures vs. LCD inferred temperatures from cultures and lake sediments (Eq. 2), respectively. Light blue lines indicate the relationships between the fractional abundance of LCDs (a–d), LDI (e) and equation 2-inferred temperatures (f) in marine surface sediments and sea surface temperature as reported by Rampen et al. (2012). The red encircled lake sample points in (e) and (f) indicate an outlier sample from Lago di Albano.

temperature when using the different monthly mean air temperatures instead of MAAT.

As an alternative approach, we used branched GDGT distributions of the lake sediments previously reported by

Blaga et al. (2010) to reconstruct summer lake temperature by application of the calibration of Pearson et al. (2011). It is likely that the algae in the lake are growing predominantly in summer and, in this way, the calibration becomes

independent from the data of the weather stations, which, as explained, have their limitations. Interestingly, the use of branched GDGT-derived temperatures results in an improved correlation of temperature with LDI-values ( $R^2 = 0.51$ ,  $p$ -value  $< 0.001$ ,  $n = 55$ , Fig. 6e). The data point from Lago di Albano (branched GDGT-derived temperature of 10.6 and an LDI value of 0.95) was identified as an outlier in the lake dataset (studentized residual = 5.027). Although the properties of Lago di Albano are not clearly different from other lakes in this study (Blaga et al., 2010), the LCD fraction is characterized by an unusual dominance of  $C_{30}$  1,15-alkyl diol (>50%) and a corresponding high LDI value, not observed in any other lake. Exclusion of this data point strongly improves the correlation with LDI, resulting in an  $R^2$  value of 0.64 ( $n = 54$ ). Comparison with the marine LDI-temperature correlation shows a difference in slope (Fig. 6e). This may be partially explained by the fact that the LDI calibration for marine environments is based on annual mean sea surface temperatures, while the LDI calibration in lakes is based on branched GDGT-derived summer temperatures.

We also applied a multiple linear regression on relative abundances of the four most dominant LCDs, e.g.  $C_{28}$  and  $C_{30}$  1,13-alkyl diol and  $C_{30}$  and  $C_{32}$  1,15-alkyl diol and branched GDGT derived summer temperatures. Again, data from Lago di Albano is identified as an outlier (studentized residual = 4.577) and when excluding this data point, the following model is obtained (Fig. 6f):

$$\begin{aligned} \text{Temperature} (^{\circ}\text{C}) = & 26.8 - 25.9 * \text{RA}_{C_{28},1,13\text{-alkyl diol}} \\ & - 54.3 * \text{RA}_{C_{30},1,13\text{-alkyl diol}} + 7.4 \\ & * \text{RA}_{C_{30},1,15\text{-alkyl diol}} \end{aligned} \quad (2)$$

The correlation observed for this model ( $R^2 = 0.74$ ,  $p$ -value  $< 0.001$ ,  $n = 54$ ) does suggest that also in lacustrine environments, LCD distributions are affected by temperature and may therefore also have potential for lacustrine temperature reconstruction. The improved correlation with proxy temperature rather than measured annual mean temperatures may be that the proxy temperature is more representative of the lake temperature during production of the LCDs. Nevertheless, the scatter remains substantial, and may have several causes. First of all, the branched GDGT-derived temperature has a much larger error than instrumentally measured temperatures that are typically used for proxy calibration. In addition, the source organisms for the lipids used in both proxies may bloom at different times or depths, thereby registering different temperatures. The scatter may also be due to the occurrence of different eustigmatophyte species in the different lakes, which may have different blooming periods, but may also possess different LCD adaptations to temperature, something also observed for alkenones in lacustrine haptophyte algae (see Castañeda and Schouten, 2011 and references cited therein).

When the obtained regression model is applied to the LCD data of the eustigmatophyte cultures analyzed in this study, the temperature values obtained for the *N. gaditana* and *V. stellata* cultures also show a positive correlation with growth temperature. Remarkably, the correlation

slopes for *N. gaditana* and *V. stellata* are not significantly different from the lake dataset (Fig. 6f,  $p$ -values of 0.502 and 0.665, respectively). In addition, although the intercept for *N. gaditana* clearly differs from the lake dataset ( $p$ -value  $< 0.001$ ), the  $y$ -intercepts for *V. stellata* and the lake dataset are not significantly different ( $p$ -value = 0.541). The similarities observed between LCD distributions in lakes and *Vischeria* species, and their correlations with temperature, may indicate that *Vischeria* species, and possibly species of the family Eustigmataceae in general, are an important source for lacustrine LCDs. Alternatively, and perhaps more likely, LCD distributions in lacustrine sources may be similarly affected by environmental conditions as in *Vischeria* species.

The results of this study show the potential of LCDs for lacustrine temperature reconstruction. However, calibration of this new proxy requires a much larger lake dataset, including well-constrained annual and seasonal lake water temperatures, but also pH, alkalinity and other factors, which may affect LCD distributions. In addition, analysis of sediment trap and suspended particulate matter samples, and new eustigmatophyte isolates may provide valuable information.

## 5. CONCLUSIONS

LCDs were detected in ten eustigmatophyte species with distinct LCD patterns for different eustigmatophyte families. Identification of LCDs in fifty-eight out of sixty-two studied lake sediments indicates that these lipids are common biomarkers in lacustrine environments. Comparison of our results with previously published LCD distributions in marine sediments indicates higher relative abundances of  $C_{32}$  1,15-alkyl diol in cultures and most lake sediments, indicating different sources. From all eustigmatophyte species analyzed in this study, LCD distributions from the algal family Eustigmataceae are most similar to the LCD distributions of aquatic environments. Cultivation of several species at a range of temperatures show an increase in saturated  $C_{32}$  1,15-alkyl diol and a corresponding decrease in mono-unsaturated  $C_{32}$  1,15-alkyl diol with increasing temperature in *N. gaditana* and *G. sculpta*. In natural environments, however, the mono-unsaturated  $C_{32}$  1,15-alkyl diol is rarely abundant, whereas relative abundances of  $C_{28}$  and  $C_{30}$  1,13- and  $C_{30}$  1,15-alkyl diols in marine environments do correlate with temperature. Relative abundances of these individual LCDs in lakes show no correlation with mean annual air temperature. However, the Long chain Diol Index, (LDI) based on these three lipids, does correlate with temperature, and an even stronger correlation is observed using a multiple linear regression. The strongest correlation was obtained by performing multiple linear regression on relative abundances of  $C_{28}$  and  $C_{30}$  1,13- and  $C_{30}$  1,15-alkyl diols and branched GDGT-derived temperatures from the same lakes. The  $R^2$  value of 0.74 obtained for this model suggests that LCDs may also be applicable as paleotemperature proxies for lacustrine environments. However, more validation studies are needed before this proxy can be applied with confidence.

## ACKNOWLEDGEMENTS

We thank J.J. Brocks, J.K. Volkman and two anonymous reviewers for their constructive comments. The research leading to these results has been funded from a VENI Grant to S.W.R. and Grant ZKO-VTOP/010-17 to J.S.S.D., provided by the Earth and Life Sciences Division of the Netherlands Organization for Scientific Research (NWO-ALW), and it has received funding from the European Research Council (ERC) under the European Union's Seventh Framework Program (FP7/2007-2013) ERC grant agreement [339206] to S.S.

## APPENDIX A. SUPPLEMENTARY DATA

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.gca.2014.08.033>.

## REFERENCES

- Blaga C. I., Reichart G.-J., Schouten S., Lotter A. F., Werne J. P., Kosten S., Mazzeo N., Lacerot G. and Sinninghe Damsté J. S. (2010) Branched glycerol dialkyl glycerol tetraethers in lake sediments: can they be used as temperature and pH proxies? *Org. Geochem.* **41**, 1225–1234.
- Britton T., Anderson C. L., Jacquet D., Lundqvist S. and Bremer K. (2007) Estimating divergence times in large phylogenetic trees. *Syst. Biol.* **56**, 741–752.
- Buser H.-R., Arn P., Guerin P. and Rauscher S. (1983) Determination on double bond position in mono-unsaturated acetates by mass spectrometry of dimethyl disulfide adducts. *Anal. Chem.* **55**, 818–822.
- Castañeda I. S. and Schouten S. (2011) A review of molecular organic proxies for examining modern and ancient lacustrine environments. *Quat. Sci. Rev.* **30**, 2851–2891.
- De Leeuw J. W., Rijpstra W. I. C. and Schenck P. A. (1981) The occurrence and identification of C<sub>30</sub>, C<sub>31</sub> and C<sub>32</sub> alkan-1,15-diols and alkan-15-one-1-ols in Unit I and Unit II Black Sea sediments. *Geochim. Cosmochim. Acta* **45**, 2281–2285.
- Déry S. J., Mlynowski T. J., Hernández-Henríquez M. A. and Straneo F. (2011) Interannual variability and interdecadal trends in Hudson Bay streamflow. *J. Marine Syst.* **88**, 341–351.
- Fawley K. P., Eliaš M. and Fawley M. W. (in press) The diversity and phylogeny of the commercially important algal class Eustigmatophyceae, including the new clade *Goniochloridales*. *J. Appl. Phycol.* DOI 10.1007/s10811-013-0216-z.
- Kimura M. (1980) A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J. Mol. Evol.* **16**, 111–120.
- KNMI, 1998. Royal Netherlands Meteorological Institute (KNMI), World Climate Information (WKI 2.0), available online at: <http://www.knmi.nl/klimatologie/normalen1971-2000/wki.html>.
- Livingstone D. M. and Lotter A. F. (1998) The relationship between air and water temperatures in lakes of the Swiss Plateau: a case study with palaeolimnological implications. *J. Paleolimnol.* **19**, 181–198.
- Loomis S. E., Russell J. M., Ladd B., Street-Perrott F. A. and Sinninghe Damsté J. S. (2012) Calibration and application of the branched GDGT temperature proxy on East African lake sediments. *Earth Planet. Sci. Lett.* **357–358**, 277–288.
- Lopes dos Santos R. A., Spooner M. I., Barrows T. T., De Deckker P., Sinninghe Damsté J. S. and Schouten S. (2013) Comparison of organic (U<sup>K37</sup>, TEX<sup>H86</sup>, LDI) and faunal proxies (foraminiferal assemblages) for reconstruction of late quaternary sea surface temperature variability from offshore southeastern Australia. *Paleoceanography* **28**, 377–387.
- Méjanelle L., Sanchez-Gargallo A., Bentaieb I. and Grimalt J. O. (2003) Long chain n-alkyl diols, hydroxy ketones and sterols in a marine eustigmatophyte, *Nannochloropsis gaditana*, and in *Brachionus plicatilis* feeding on the algae. *Org. Geochem.* **34**, 527–538.
- Pearson E. J., Juggins S., Talbot H. M., Weckstrom J., Rosen P., Ryves D. B., Roberts S. J. and Schmidt R. (2011) A lacustrine GDGT-temperature calibration from the Scandinavian Arctic to Antarctic: renewed potential for the application of GDGT-paleothermometry in lakes. *Geochim. Cosmochim. Acta* **75**, 6225–6238.
- Prior S. W., Fawley M. W. and Fawley K. P. (2009) DNA sequence analysis of freshwater Eustigmatophyceae, a potential source of essential fatty acids. *J. Ark. Acad. Sci.* **63**, 139–144.
- Rampen S. W., Schouten S., Koning E., Brummer G.-J. A. and Sinninghe Damsté J. S. (2008) A 90 kyr upwelling record from the northwestern Indian Ocean using a novel long-chain diol index. *Earth Planet. Sci. Lett.* **276**, 207–213.
- Rampen S. W., Schouten S., Schefuß E. and Sinninghe Damsté J. S. (2009) Impact of temperature on long chain diol and mid-chain hydroxy methyl alkanoate composition in *Proboscia* diatoms: results from culture and field studies. *Org. Geochem.* **40**, 1124–1131.
- Rampen S. W., Schouten S. and Sinninghe Damsté J. S. (2011) Occurrence of long chain 1,14 diols in *Apedinella radians*. *Org. Geochem.* **42**, 572–574.
- Rampen S. W., Schouten S., Wakeham S. G. and Sinninghe Damsté J. S. (2007) Seasonal and spatial variation in the sources and fluxes of long chain diols and mid-chain hydroxy methyl alkanoates in the Arabian Sea. *Org. Geochem.* **38**, 165–179.
- Rampen S. W., Willmott V., Kim J.-H., Uliana E., Mollenhauer G., Schefuß E., Sinninghe Damsté J. S. and Schouten S. (2012) Long chain 1,13- and 1,15-diols as a potential proxy for palaeotemperature reconstruction. *Geochim. Cosmochim. Acta* **84**, 204–216.
- Rodrigo-Gámiz M., Martínez-Ruiz F., Rampen S. W., Schouten S. and Sinninghe Damsté J. S. (2014) Sea surface temperature variations in the western Mediterranean Sea over the last 20 kyr: a dual-organic proxy (U<sup>K57</sup> and LDI) approach. *Paleoceanography* **29**, 87–98.
- Romero-Viana L., Kienel U. and Sachse D. (2012) Lipid biomarker signatures in a hypersaline lake on Isabel Island (Eastern Pacific) as a proxy for past rainfall anomaly (1942–2006 AD). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **350–352**, 49–61.
- Romero-Viana L., Kienel U., Wilkes H. and Sachse D. (2013) Growth-dependent hydrogen isotopic fractionation of algal lipid biomarkers in hypersaline Isabel Lake (Mexico). *Geochim. Cosmochim. Acta* **106**, 490–500.
- Russell N. J. and Fukunaga N. (1990) A comparison of thermal adaptation of membrane-lipids in psychrophilic and thermophilic bacteria. *FEMS Microbiol. Rev.* **75**, 171–182.
- Saitou N. and Nei M. (1987) The Neighbor-Joining method – a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* **4**, 406–425.
- Seki O., Schmidt D. N., Schouten S., Hopmans E. C., Sinninghe Damsté J. S. and Pancost R. D. (2012) Paleoclimatographic changes in the Eastern Equatorial Pacific over the last 10 Myr. *Paleoceanography* **27**, PA3224.
- Shimokawara M., Nishimura M., Matsuda T., Akiyama N. and Kawai T. (2010) Bound forms, compositional features, major sources and diagenesis of long chain, alkyl mid-chain diols in

- Lake Baikal sediments over the past 28,000 years. *Org. Geochem.* **41**, 753–766.
- Sinninghe Damsté J. S., Rampen S., Rijpstra W. I. C., Abbas B., Muyzer G. and Schouten S. (2003) A diatomaceous origin for long-chain diols and mid-chain hydroxy methyl alkanooates widely occurring in Quaternary marine sediments: Indicators for high nutrient conditions. *Geochim. Cosmochim. Acta* **67**, 1339–1348.
- Speelman E. N., Reichart G.-J., De Leeuw J. W., Rijpstra W. I. C. and Sinninghe Damsté J. S. (2009) Biomarker lipids of the freshwater fern *Azolla* and its fossil counterpart from the Eocene Arctic Ocean. *Org. Geochem.* **40**, 628–637.
- Stone P. H. and Carlson J. H. (1979) Atmospheric lapse rate regimes and their parameterization. *J. Atmos. Sci.* **36**, 415–423.
- Suutari M. and Laakso S. (1994) Microbial fatty acids and thermal adaptation. *Crit. Rev. Microbiol.* **20**, 285–328.
- Tierney J. E., Russell J. M., Eggermont H., Hopmans E. C., Verschuren D. and Sinninghe Damsté J. S. (2010) Environmental controls on branched tetraether lipid distributions in tropical East African lake sediments. *Geochim. Cosmochim. Acta* **74**, 4902–4918.
- Versteegh G. J. M., Bosch H. J. and De Leeuw J. W. (1997) Potential palaeoenvironmental information of C<sub>24</sub> to C<sub>36</sub> mid-chain diols, keto-ols and mid-chain hydroxy fatty acids; a critical review. *Org. Geochem.* **27**, 1–13.
- Villanueva L., Besseling M., Rodrigo-Gámiz M., Rampen S. W., Verschuren D. and Sinninghe Damsté J. S. (2014) Potential biological sources of long chain alkyl diols in a lacustrine system. *Org. Geochem.* **68**, 27–30.
- Volkman J. K., Barrett S. M. and Blackburn S. I. (1999) Eustigmatophyte microalgae are potential sources of C<sub>29</sub> sterols, C<sub>22</sub>–C<sub>28</sub> *n*-alcohols and C<sub>28</sub>–C<sub>32</sub> *n*-alkyl diols in freshwater environments. *Org. Geochem.* **30**, 307–318.
- Volkman J. K., Barrett S. M., Dunstan G. A. and Jeffrey S. W. (1992) C<sub>30</sub>–C<sub>32</sub> alkyl diols and unsaturated alcohols in microalgae of the class Eustigmatophyceae. *Org. Geochem.* **18**, 131–138.
- Willmott V., Rampen S. W., Domack E., Canals M., Sinninghe Damsté J. S. and Schouten S. (2010) Holocene changes in *Proboscia* diatom productivity in shelf waters of the north-western Antarctic Peninsula. *Antarct. Sci.* **22**, 3–10.

Associate editor: Jochen J. Brocks